

CONTENTS

	Page
Declaration	ii
Approval	iii
Abstract	iv
Acknowledgements	v
Contents	vi
List of tables	ix
List of figures	x
List of plates	xii
CHAPTER 1. INTRODUCTION	1
1.1 Classification	1
1.2 Shell characteristics	3
1.3 Bivalve distribution	4
1.4 Review of factors influencing the distribution of bivalves	5
1.4.1 Salinity and temperature	6
1.4.2 Nature of the substratum	9
1.4.3 Submerged benthic vegetation cover	11
1.5 Bivalve fisheries and aquaculture	12
1.6 Study area and objectives	24

CHAPTER 2. MATERIALS AND METHODS	27
2.1 Selection of sampling sites	27
2.2 Environmental factors	27
2.2.1 Temperature	28
2.2.2 Salinity	28
2.2.3 Soil texture and soil organic matter content	29
2.2.4 Seagrass coverage	29
2.3 Distribution of bivalves	30
2.4 Bivalve fishery	31
2.4.1 Estimation of total catch	31
2.4.2 The length weight relationship	31
2.5 Growth and mortality parameters	32
2.5.1 Estimation of growth parameters	32
2.5.1.1 Growth performance	33
2.5.2 Estimation of total mortality	34
2.5.3 Estimation of natural mortality	35
2.5.4 Estimation of fishing mortality	36
2.5.5 Estimation of probabilities of capture and mean length at 50% selection	36
2.6 Recruitment pattern	36
2.7 Relative Yield per Recruit	37

CHAPTER 3. RESULTS	38
3.1 Temperature	38
3.2 Salinity	38
3.3 Soil texture and organic matter content	42
3.4 Seagrass cover	42
3.5 Distribution of bivalves and factors influencing their distribution	43
3.6 Bivalve fishery	45
3.7 Length weight relationship	61
3.8 Growth and mortality	61
3.8.1 Growth	61
3.8.1.1 Growth performance	63
3.8.2 Mortality	63
3.8.2.1 Total mortality	63
3.8.2.2 Natural mortality	74
3.8.2.3 Fishing mortality rates	76
3.9 Exploitation rates	76
3.10 Probabilities of capture and mean size at first capture	76
3.11 Recruitment pattern	80
3.12 Relative yield per recruit	80
CHAPTER 4. DISCUSSION	90
5. REFERENCES	104

LIST OF TABLES

Table No.		Page
1.	Classification of the three bivalve species studied	2
2.	List of major species of clams, cockles and arkshells that are harvested	15-20
3.	Annual landings for major clam/ cockle (mt) fisheries of the world from 1987 to 1993	22
4.	Mean values for Air and Water temperature of sampling sites in Dutch bay and Puttalam lagoon during the study period	39
5.	Mean salinity values (ppt) at sampling sites during the study period	41
6.	The values for correlation coefficient between abundance of bivalves and some environmental parameters	44
7.	The monthly production of the three species of bivalves and the income earned by the fishermen in Dutch Bay during 1992/1993 fishing seasons	55
8.	Length-weight relationship for the three bivalve species in Puttalam lagoon and Dutch Bay	62
9.	Growth parameter estimates for three bivalve species in Puttalam lagoon and Dutch Bay	64
10.	Estimates for mortality rates, exploitation rates(E) and sizes of first capture (L_c) of <i>G. tumidum</i> (GT), <i>M. hiantina</i> (MH) and <i>M. opima</i> (MO) in Dutch Bay area	64

LIST OF FIGURES

Figure no:	Page
1. Location of sampling sites, fishing villages and major fishing grounds in Puttalam lagoon and Dutch Bay	25
2. Variation of bivalve abundance with different environmental parameters	40
3. Monthly production of <i>G. tumidum</i> (with shell) in Dutch Bay	56
4. Monthly production of <i>M. hiantina</i> (with shell) in Dutch Bay	57
5. Monthly production of <i>M. opima</i> (with shell) in Dutch Bay	58
6. The average monthly income/fisher during fishing season	59
7. Variation of bivalve production with fishing effort	60
8. Restructured and percentage length frequency data and superimposed growth curves for <i>Gafrarium tumidum</i> in Puttalam lagoon	65
9. Restructured and percentage length frequency data and superimposed growth curves for <i>M. hiantina</i> in Puttalam lagoon	66
10. Restructured and percentage length frequency data and superimposed growth curves for <i>M. opima</i> in Puttalam lagoon	67
11. Restructured and percentage length frequency data and superimposed growth curves for <i>G. tumidum</i> in Dutch Bay	68

12.	Restructured and percentage length frequency data and superimposed growth curves for <i>M. hiantina</i> in Dutch Bay	69
13.	Restructured and percentage length frequency data and superimposed growth curves for <i>M. opima</i> in Dutch Bay	70
14.	Length -converted catch curves for <i>G. tumidum</i>	71
15.	Length -converted catch curves for <i>M. hiantina</i>	72
16.	Length -converted catch curves for <i>M. opima</i>	73
17.	Time lag cross correlation analysis for the three bivalve species	75
18.	The selection curves for <i>G. tumidum</i> in Dutch Bay	77
19.	The selection curves for <i>M. hiantina</i> in Dutch Bay	78
20.	The selection curves for <i>M. opima</i> in Dutch Bay	79
21.	Recruitment pattern of <i>G. tumidum</i> in Puttalam lagoon	81
22.	Recruitment pattern of <i>G. tumidum</i> in Dutch Bay	82
23.	Recruitment pattern of <i>M. hiantina</i> in Puttalam lagoon	83
24.	Recruitment pattern of <i>M. hiantina</i> in Dutch Bay	84
25.	Recruitment pattern of <i>M. opima</i> in Puttalam lagoon	85
26.	Recruitment pattern of <i>M. opima</i> in Dutch Bay	86
27.	Relative yield per recruit values as functions of exploitation rates for <i>G. tumidum</i>	87
28.	Relative yield per recruit values as functions of exploitation rates for <i>M. hiantina</i>	88
29.	Relative yield per recruit values as functions of exploitation rates for <i>M. opima</i>	89

LIST OF PLATES

Plate No.		Page
1.	Bivalve beds exposed during the low tide period in Dutch Bay area.	46
2.	Clams and cockles buried just below the surface of the soil.	47
3.	Bivalves collected from Dutch bay area and window-pane oyster shells that are used to collect bivalves.	48
4.	Harvesting bivalves in Dutch Bay area using the shells of window-pane oyster (<i>Placuna placenta</i>).	49
5.	Harvesting bivalves in Dutch Bay area during the high tide period.	50
6.	Harvesting bivalves in Dutch Bay area during the low tide period.	51
7.	<i>Gafrarium tumidum</i> collected from Dutch Bay area.	52
8.	Bivalve shells used for lime production.	54