

CONTENTS

	Page No
I. LIST OF TABLES	vii
II. LIST OF FIGURES	x
III. LIST OF APPENDICES	xii
IV. LIST OF PLATES	xii
V. ACKNOWLEDGEMENTS	xiii
VI. ABSTRACT	xv
1. INTRODUCTION	1
2. LITERATURE REVIEW	4
2.1. Composition of Fish Lipids	4
2.1.1. Fatty acids	5
2.1.2. Neutral lipids	8
2.1.3. Polar lipids	10
2.2. Distribution of Lipids in Fish	12
2.3. Extraction Modification and Utilization of Fish Lipids	14
2.3.1. Extraction of fish lipids	14
2.3.2. Modification of fish lipids	15
2.3.3. Utilization of fish lipids	16
2.4. Analysis of Composition and Quality of Fish Lipids	18
2.4.1. Isolation and purification of fish lipids	18
2.4.2. Determination of lipid classes and fatty acid profile	20
2.4.3. Determination of quality of the fish lipids	25

2.5.	Nutritional and Pharmaceutical Importance of Fish Lipids	27
2.5.1.	Fatty acids	27
2.5.2.	Cholesterol	31
2.5.3.	Vitamins	32
2.5.4.	Calorific value	32
2.6.	Changes in the Lipids	32
2.6.1.	Lipid autoxidation	33
2.6.2.	Photooxygenation	35
2.6.3.	Lipid autolysis	39
2.6.4.	Interaction between lipids and proteins	41
2.7.	Preservation of Lipids	42
2.7.1.	Storage at low temperature	43
2.7.2.	Packaging	43
2.7.3.	Uses of antioxidants in preservation of fish lipids	44
3.	EXPERIMENTAL	52
3.1.	Proximate Composition of Fish	52
3.1.1.	Determination of moisture	52
3.1.1.1.	Materials	52
3.1.1.2.	Method	52
3.1.2.	Determination of ash	53
3.1.2.1.	Materials	53
3.1.2.2.	Method	53
3.1.3.	Determination of lipid	54
3.1.3.1.	Materials	54
3.1.3.2.	Method	54
3.1.4.	Determination of protein	55
3.1.4.1.	Materials	55
3.1.4.2.	Method	56

3.2.	Extraction and Quality Determination of Fish Lipids	57
3.2.1.	Extraction of fish lipids	57
3.2.1.1.	Extraction by wet rendering	57
3.2.1.1.1.	Materials	57
3.2.1.1.2.	Method	58
3.2.1.2.	Extraction by steaming	58
3.2.1.2.1.	Materials	58
3.2.1.2.2.	Method	58
3.2.1.3.	Acid silage extraction	58
3.2.1.3.1.	Materials	58
3.2.1.3.2.	Method	59
3.2.1.4.	Microbial silage extraction	59
3.2.1.4.1.	Materials	59
3.2.1.4.2.	Method	59
3.2.1.5.	Bligh & Dyer extraction	60
3.2.1.5.1.	Materials	60
3.2.1.5.2.	Method	60
3.2.2.	Determination of quality of extracted fish lipids	60
3.2.2.1.	Determination of peroxide value	60
3.2.2.1.1.	Materials	60
3.2.2.1.2.	Method	61
3.2.2.2.	Determination of free fatty acid value	62
3.2.2.2.1.	Materials	62
3.2.2.2.2.	Method	62
3.2.2.3.	Determination of iodine value	63
3.2.2.3.1.	Materials	63
3.2.2.3.2.	Method	63
3.2.2.4.	Determination of fatty acid composition	64
3.2.2.4.1.	Materials	64
3.2.2.4.2.	Method	65

3.3.	Lipid and Fatty Acid Composition of Selected Fish	67
3.3.1.	Materials	67
3.3.2.	Method	67
3.4.	Fatty Acid Composition (Total) of Small Pelagics	69
3.4.1.	Materials	69
3.4.2.	Method	69
3.5.	Seasonal Variation of Quality and Quantity of Fish Lipids in Selected Species	70
3.5.1.	Materials	70
3.5.2.	Method	70
3.6.	Quality Changes of Fish and Fish Lipids During Storage	71
3.6.1.	Storage of fish and fish lipids at room temperature ($32^{\circ}\pm 1\text{C}$)	71
3.6.1.1.	Materials	71
3.6.1.2.	Method	71
3.6.2.	Storage of fish in ice ($0^{\circ}\pm 1\text{ C}$)	72
3.6.2.1.	Materials	72
3.6.2.2.	Method	72
3.6.3.	Storage of fish and fish lipids in freezer ($-18^{\circ}\pm 1\text{ C}$)	73
3.6.3.1.	Materials	73
3.6.3.2.	Method	73
3.6.4.	Determination of total volatile bases (TVB) by Conway-Byrnes method	74
3.6.4.1.	Materials	74
3.6.4.2.	Method	75
3.7.	Preservation of Fish Lipids Using Natural Anti-oxidants	76
3.7.1.	Studies on the efficiency of some natural plant product extracts in the preservation of fish lipids.	76
3.7.1.1.	Materials	76
3.7.1.2.	Methods	77

3.7.2.	Quantification of preservative action of crude Indian gooseberry extract	78
3.7.2.1.	Materials	78
3.7.2.1.	Method	78
3.7.3.	Evaluation of anti-oxidant activity of Indian gooseberry using different solvent systems	79
3.7.3.1.	Materials	79
3.7.3.2.	Method	79
3.7.4.	Purification of crude Indian gooseberry extract and the examination of preservative action of purified fractions	80
3.7.4.1.	Materials	80
3.7.4.2.	Method	81
4.	RESULTS AND DISCUSSION	83
4.1.	Proximate Composition of Fish	83
4.1.1	Moisture content	83
4.1.2.	Ash content	83
4.1.3.	Lipid content	84
4.1.4.	Protein content	85
4.2.	Extraction and Quality Determination of Fish Lipids	88
4.3.	Lipid and Fatty Acid Composition of Selected Fish	94
4.3.1.	Lipid composition of selected fish species	94
4.3.2.	Fatty acid composition of lipid classes of White sardinella	96
4.4.	Fatty Acid Composition of Small Pelagics	98
4.4.1.	Saturated fatty acids	98
4.4.2.	Monounsaturated fatty acids	99
4.4.3.	Polyunsaturated fatty acids	99
4.5.	Seasonal Variation of Quality and Quantity of Fish Lipids in Selected Species	108
4.5.1.	Variation of lipid content	108

4.5.2.	Variation of fatty acids	110
4.6.	Quality Changes of Fish and Fish Lipids During Storage	115
4.6.1.	Changes observed in the quality of fish and fish lipids during storage at room temperature ($32^{\circ} \pm 1\text{C}$)	115
4.6.2.	Changes observed in the quality of fish and lipids during ice storage ($0^{\circ} \pm 1\text{C}$)	118
4.6.3.	Changes observed in the quality of fish and fish lipids during storage at $-18^{\circ} \pm 1\text{C}$	120
4.7.	Preservation of Fish Lipids Using Natural Anti-oxidants	126
4.7.1.	Studies on the effect of some natural plant extracts in the preservation of fish lipids	126
4.7.2	Quantification of preservative action of crude Indian Gooseberry extract	135
4.7.3.	Evaluation of anti-oxidant activity of Indian gooseberry using different solvent systems to extract active components	141
4.7.4.	Investigation of antioxidant activity of purified compounds from Indian gooseberry	148
5.	CONCLUSIONS	155
	LITEATURE CITED	161
	APPENDIX	182
	PLATES	184
	RESEARCH PUBLICATIONS AND COMMUNICATIONS FROM THE STUDY	189

LIST OF TABLES

	Page No	
Table-2.1.1	Intramuscular phospholipid distribution of Sucker (<i>Catostomus commersoni</i>).	11
Table-2.1.2	Comparison between fatty acids of phospholipids and triglycerides in fish species.	11
Table-2.3.1	Lipid content of principal fish used in fish lipid production.	15
Table-2.5.1	Cholesterol content in some fish and seafood lipids.	31
Table-2.6.1	Relative rates of oxidation.	36
Table-4.1.1	Proximate composition of fish.	86
Table-4.2.1	Percentage of lipids extracted and its quality	89
Table-4.2.2	Changes of fatty acids of fish lipids during extraction	91
Table-4.3.1	Lipid composition of fish	94
Table-4.3.2	Fatty acid composition of lipid classes of White sardinella.	97
Table-4.4.1	List of species studied for fatty acid composition.	104
Table-4.4.2	Fatty acid composition of small pelagics.	105
Table-4.4.3	Content of fatty acids present in small pelagics.	106
Table-4.5.1	Seasonal variation of fatty acid composition of White sardinella	111
Table-4.5.2	Seasonal variation of fatty acid composition of <i>Thryssa.sp.</i>	112
Table-4.6.1.1	Changes observed in the quality of fish during storage at room temperature	115
Table-4.6.1.2	Changes observed in the quality of extracted fish lipids during storage at room temperature	116

Table-4.6.2.1	Changes observed in the quality of fish and fish lipids during ice storage.	118
Table-4.6.3.1	Changes of peroxides, free fatty acids and total volatile nitrogen of fish and fish lipids during storage at -18°C.	121
Table-4.6.3.2	Variation of fatty acid composition of fish during storage at-18°C.	124
Table-4.6.3.3	Variation of fatty acid composition of extracted fish lipids during storage at -18°C.	125
Table-4.7.1.1	Levels of peroxides of fish lipids treated with plant component extracts during storage at 30°C	127
Table-4.7.1.2	Levels of free fatty acids of fish lipids treated with plant component extracts during storage at 30°C	128
Table-4.7.1.3	Levels of saturated fatty acids of fish lipids treated with Indian Gooseberry (<i>P. emblica</i>) extract during storage at 30°C	130
Table-4.7.1.4	Levels of monounsaturated fatty acids of fish lipids treated with Indian Gooseberry (<i>P. emblica</i>) extract during storage at 30°C	131
Table-4.7.1.5	Levels of polyunsaturated fatty acids of fish lipids treated with Indian Gooseberry (<i>P. emblica</i>) extract during storage at 30°C	133
Table-4.7.2.1	Levels of peroxides of fish lipids treated with different concentrations of Indian gooseberry during storage at 30°C	136
Table-4.7.2.2	Levels of free fatty acids of fish lipids treated with different concentrations of Indian gooseberry during storage at 30°C	137
Table-4.7.2.3	Levels of saturated fatty acids of fish lipids treated with different concentrations of Indian gooseberry during storage at 30°C	138
Table-4.7.2.4	Levels of monounsaturated fatty acids of fish lipids treated with different concentrations of Indian gooseberry during storage at 30°C	139
Table-4.7.2.5	Levels of polyunsaturated fatty acids of fish lipids treated with different concentrations of Indian gooseberry during storage at 30°C	140
Table-4.7.3.1	Levels of peroxides in fish lipids during storage with different solvent extracts of Indian Gooseberry.	142

Table-4.7.3.2	Levels of free fatty acids in fish lipids during storage with different solvent extracts of Indian Gooseberry.	143
Table-4.7.3.3	Variation of saturated fatty acids of fish lipids treated with different solvent extracts of Indian gooseberry during storage at 30°C	144
Table-4.7.3.4	Variation of monounsaturated fatty acids of fish lipids treated with different solvent extracts of Indian gooseberry during storage at 30°C	145
Table-4.7.3.5	Variation of polyunsaturated fatty acids of fish lipids treated with different solvent extracts of Indian gooseberry during storage at 30°C	146
Table-4.7.4.1	Levels of peroxides in fish lipids during storage with fractionated Indian gooseberry	148
Table-4.7.4.2	Levels of free fatty acids in fish lipids during storage with fractionated Indian gooseberry	150
Table-4.7.4.3	Levels of saturated fatty acids in fish lipids during storage with fractionated Indian gooseberry	151
Table-4.7.4.4	Levels of monounsaturated fatty acids in fish lipids during storage with fractionated Indian gooseberry	152
Table-4.7.4.5	Levels of polyunsaturated fatty acids in fish lipids during storage with fractionated Indian gooseberry	153

LIST OF FIGURES

	Page No
Figure-2.1.1 Triacylglycerol (Triglyceride) of fish lipids.	4
Figure-2.2.1 Distribution of total fat in various body parts and organs of mackerel (<i>S. scombrus</i>) and capelin (<i>Mallotus villosus</i>) of Norwegian origin.	13
Figure-2.5.1 Key steps in conversion of dietary linoleic acid (18:2 n-6) to arachidonic acid (20:4 n-6) and thence to prostanoids and leukotrienes	28
Figure-2.7.1 Chemical structures of the most commonly used synthetic antioxidants	46
Figure-2.7.2 Chemical structures of a selection of natural compounds with antioxidant properties	48
Figure-3.2.1 Temperature variation of the column used in the fatty acid analysis.	66
Figure-4.1.1 Lipid content of fish	87
Figure-4.4.1 The relationship between lipid content (%) and n-3 polyunsaturated fatty acids (%)	103
Figure-4.5.1 Seasonal variation of lipid content in White sardinella	109
Figure-4.5.2 Seasonal variation of lipid content in <i>Thryssa sp</i>	109
Figure-4.5.3 Seasonal variation of lipid content in Streaked spinefoot	109
Figure-4.5.4 Seasonal variation of lipid content in Silver belly	109
Figure-4.7.1.1 Variation of peroxides of fish lipids treated with plant component extracts during storage at 30°C.	127
Figure-4.7.1.2 Variation of free fatty acids of fish lipids treated with plant component extracts during storage at 30°C	129
Figure-4.7.1.3 Variation of saturated fatty acids of fish lipids treated with Indian gooseberry extract during storage at 30°C	130
Figure-4.7.1.4 Variation of monounsaturated fatty acids of fish lipids treated with Indian gooseberry extract during storage at 30°C	131
Figure-4.7.1.5 Variation of polyunsaturated fatty acids of fish lipids treated with Indian gooseberry extract during storage at 30°C	133

Figure-4.7.2.1	Variation of peroxides of fish lipids treated with different concentrations of Indian gooseberry during storage at 30°C	136
Figure-4.7.2.2	Variation of free fatty acids of fish lipids treated with different concentrations of Indian gooseberry during storage at 30°C	137
Figure-4.7.2.3	Variation of saturated fatty acids of fish lipids treated with different concentrations of Indian gooseberry during storage at 30°C	138
Figure-4.7.2.4	Variation of monounsaturated fatty acids of fish lipids treated with different concentrations of Indian gooseberry during storage at 30°C	139
Figure-4.7.2.5	Variation of polyunsaturated fatty acids of fish lipids treated with different concentrations of Indian gooseberry during storage at 30°C	140
Figure-4.7.3.1	Variation of peroxides of fish lipids treated with different solvent extracts of Indian gooseberry during storage at 30°C	142
Figure-4.7.3.2	Variation of free fatty acids of fish lipids treated with different solvent extracts of Indian gooseberry during storage at 30°C	143
Figure-4.7.3.3	Variation of saturated fatty acids of fish lipids treated with different solvent extracts of Indian gooseberry during storage at 30°C	144
Figure-4.7.3.4	Variation of monounsaturated fatty acids of fish lipids treated with different solvent extracts of Indian gooseberry during storage at 30°C	145
Figure-4.7.3.5	Variation of polyunsaturated fatty acids of fish lipids treated with different solvent extracts of Indian gooseberry during storage at 30°C	147
Figure-4.7.4.1	Changes in the levels of peroxides in fish lipids during storage with fractionated Indian gooseberry	149
Figure-4.7.4.2	Changes in the levels of free fatty acids in fish lipids during storage with fractionated Indian gooseberry	150
Figure-4.7.4.3	Changes in the levels of saturated fatty acids in fish lipids during storage with fractionated Indian gooseberry	151
Figure-4.7.4.4	Changes in the levels of monounsaturated fatty acids in fish lipids during storage with fractionated Indian gooseberry	152
Figure-4.7.4.5	Changes in the levels of polyunsaturated fatty acids in fish lipids during storage with fractionated Indian gooseberry	153

LIST OF APPENDICES

	Page No
Appendix –1 Chilaw and Negombo fish landing sites in North-Western coast Of Sri Lanka.	182
Appendix –2 Calibration curve for lipid class analysis	183

LIST OF PLATES

	Page No
Plate – 1 Gas chromatograph (Shimadzu GC-14A)	184
Plate – 2 Lipids of White sardinella, <i>Thryssa .sp</i> , Streaked spinefoot and Silver belly developed on TLC plates.	185
Plate – 3 Fish species selected for seasonal variation studies	186
Plate – 4 Fruit of Indian Gooseberry	187
Plate – 5 Soxhlet extraction of Indian gooseberry	188