

# CONTENTS

	Page No.
Declaration	ii
Approval of the Head of the Department	iii
Abstract	iv
Acknowledgements	vi
List of Tables	xiii
List of Figures	xvi
List of Plates	xviii
<b>1. INTRODUCTION</b>	<b>1</b>
1.1 Shrimp culture industry	1
1.1.1 Shrimp culture industry - Global status	1
1.1.2 Shrimp culture industry - Sri Lankan status	5
1.2 Quality control of fishery products	10
1.3 Heavy metal contamination in the aquatic environment	14
1.4 Public health significance of lead, cadmium & chromium	18
1.5 Hypothesis tested, Aims of the Study and Specific Objectives	22
<b>2. MATERIALS &amp; METHODS</b>	<b>24</b>
2.1 Water sources & sampling sites	24
2.2 Sampling of water, sediment & shrimp	24

2.2.1	Water	24
2.2.2	Sediment	25
2.2.3	Shrimp	25
2.3	Shrimp feed	26
2.4	Processing of samples before digestion	26
2.4.1	Sediment	26
2.4.2	Shrimp	26
2.4.3	Shrimp feed	27
2.5	Digestion of sediment	27
2.6	Digestion of shrimp tissue & feed	27
2.7	Analysis of lead, cadmium and chromium	28
2.8	Bio-transfer factors	31
2.9	Statistical Analysis	32
<b>3.</b>	<b>RESULTS</b>	<b>40</b>
3.1	Levels of lead, chromium and cadmium in the water and sediments of four water sources that fed the shrimp farms studied	40
3.1.1	Puttalam Lagoon	40
3.1.2	Mundal Lagoon	41
3.1.3	Gembarandiya Lagoon	41
3.1.4	Dutch Canal	42
3.1.5	Overall comparison of the metal levels in the water and sediments of the four water sources	43

3.2	Levels of lead, cadmium and chromium in the sediments of the shrimp farms studied.	44
3.2.1	Shrimp farms fed from Puttalam Lagoon	44
3.2.2	Shrimp farms fed from Mundal Lagoon	44
3.2.3	Shrimp farms fed from Gembarandiya Lagoon	45
3.2.4	Shrimp farms fed from Dutch Canal	45
3.2.5	Water source wise comparison of the levels of lead, cadmium and chromium in farm sediments fed from four water sources	46
3.3	Levels of lead, chromium and cadmium in the shrimp, <i>Penaeus monodon</i> cultured in water from four water sources	47
3.3.1	Shrimp cultured in water from Puttalam Lagoon	47
3.3.2	Shrimp cultured in water from Mundal Lagoon	47
3.3.3	Shrimp cultured in water from Gembarandiya Lagoon	48
3.3.4	Shrimp cultured in water from Dutch Canal	48
3.3.5	Water source wise comparison of metal levels in the shrimp cultured in water from four water sources	48
3.4.	Formulated shrimp feed used during the culture cycles and the levels of lead, chromium and cadmium in the shrimp feed	49
3.4.1	Formulated shrimp feed used during the culture cycles	49
3.4.2	Levels of lead, chromium and cadmium in the shrimp feed	50

3.5	Overall residue levels of lead, chromium and cadmium in shrimp tissue cultured in selected farms located in the North Western Province of Sri Lanka	51
3.6	Bio-transfer factors of the three metals in shrimp tissue	52
<b>4.</b>	<b>DISCUSSION</b>	<b>79</b>
<b>5.</b>	<b>REFERENCES</b>	<b>90</b>
	<b>APPENDICES</b>	<b>96</b>
	Appendix 01: Variation of each metal concentration ( $\mu\text{g g}^{-1}$ dry weight) in sediments among 12 farms connected to four water sources	96
	Appendix 02: Variation of each metal concentration ( $\mu\text{g g}^{-1}$ wet weight) in shrimp among 12 farms connected to four water sources	97

## List of Tables

	Page No.
Table 1.1: Global aquaculture production and shrimp production	4
Table 1.2: Country wise export of shrimp and prawn in Sri Lanka, 1996 – 2000	9
Table 1.3: Permissible levels of lead, cadmium and chromium in shellfish and finfish for human consumption	13
Table 1.4: Proposed quality standards for different use classes of coastal waters of Sri Lanka	17
Table 3.1: Comparison of the levels of lead, chromium and cadmium in water ( $\text{mg l}^{-1}$ ) and sediments ( $\mu\text{g g}^{-1}$ dry weight) of four water sources that fed the shrimp farms studied	55
Table 3.2: Comparison of the levels of lead, chromium and cadmium in water ( $\text{mg l}^{-1}$ ) and sediments ( $\mu\text{g g}^{-1}$ dry weight) among four water sources that fed the shrimp farms studied	56
Table 3.3: Variation of the levels of three metals in the sediments of shrimp farms studied	57
Table: 3.4: Comparison of the metal levels in farm sediments in three farms connected to Puttalam lagoon	58
Table 3.5: Comparison of the metal levels in farm sediments in three farms connected to Mundal lagoon	59

Table 3.6:	Comparison of the metal levels in farm sediments in three farms connected to Gembarandiya lagoon	60
Table 3.7:	Comparison of the metal levels in farm sediments in three farms connected to Dutch canal	61
Table 3.8:	Variation of the metal levels in sediments ( $\mu\text{g g}^{-1}$ dry weight) of the farms fed from four water sources studied	62
Table 3.9:	Variation among the levels of lead, chromium and cadmium in tiger shrimp, <i>Penaeus monodon</i> cultured in twelve farms	64
Table 3.10:	Comparison of the metal levels in shrimp tissue in three farms connected to Puttalam lagoon	66
Table 3.11:	Comparison of the metal levels in shrimp tissue in the three farms connected to Mundal lagoon	67
Table 3.12:	Comparison of the metal levels in shrimp tissue in three farms connected to Gembarandiya lagoon	68
Table 3.13:	Comparison of the metal levels in shrimp tissue in three farms connected to Dutch canal	69
Table 3.14:	Comparison of the metal levels in shrimp ( $\mu\text{g g}^{-1}$ wet weight) cultured in water from the four different water sources	70
Table 3.15:	The formulated shrimp feed used by the different farms during the culture cycle	72
Table 3.16:	Levels of lead, chromium and cadmium ( $\mu\text{g g}^{-1}$ wet weight) in different types of feed used by the shrimp farms	73

Table 3.17:	Residue levels of lead, chromium and cadmium in shrimp ( <i>Penaeus monodon</i> ) cultured in selected farms of the North Western Province, Sri Lanka	75
Table 3.18:	Bio-transfer factors of lead, chromium and cadmium in relation to <i>Penaeus monodon</i> cultured in water from four water sources	76
Table 3.19:	Comparison of the transfer factors of lead, chromium and cadmium in <i>Penaeus monodon</i> in relation to water, sediments and formulated feed	77

## List of Figures

	Page No.
Figure 1.1: Shrimp Production in Sri Lanka, 1990 – 2000	8
Figure 2.1: Location of the four selected water sources that fed the shrimp farms studied	34
Figure 2.2: Location of the sampling sites in Puttalam lagoon and the three selected farms fed from this water source	37
Figure 2.3: Location of the sampling sites in Mundal lagoon and the three selected farms fed from this water source	38
Figure 2.4: Location of the sampling sites in Gembarandiya lagoon and the three selected farms fed from this water source	39
Figure 2.5: Location of the sampling sites in Dutch canal (North & South) and the three selected farms fed from this water Source	40
Figure 3.1: Comparison of the levels of lead, cadmium and chromium ( $\mu\text{g g}^{-1}$ dry weight) in the sediments of the farms among four water sources	63
Figure 3.2: Comparison of the levels of lead, cadmium and chromium in the muscle tissue of shrimp, <i>Penaeus monodon</i> cultured in three farms connected to each of the four water sources	65

Figure 3.3:	Comparison of the levels of lead, cadmium and chromium in the muscle tissue of shrimp ( $\mu\text{g g}^{-1}$ wet weight) among four water sources	71
Figure 3.4:	Comparison of the levels of lead, chromium and cadmium in the formulated feed used in the farms	74
Figure 3.5:	Comparison of the bio-transfer factors of lead, chromium and cadmium in relation to water, sediments of the water sources, sediments of the farms and formulated feed	78

## List of Plates

	Page No.
Plate 1.1: Black tiger shrimp, <i>Penaeus monodon</i>	3
Plate 2.1: Puttalam Lagoon	35
Plate 2.2: Mundal Lagoon	35
Plate 2.3: Gembarandiya Lagoon	36
Plate 2.4: Dutch Canal	36