

CONTENTS

	Page No.
1. INTRODUCTION	1
2. REVIEW OF LITERATURE	8
2.1. Accumulation of Plastic Debris on Sandy Beaches	8
2.1.1. Factors affecting plastic litter distribution on beaches	9
2.1.2. Plastic litter distribution in beaches	10
2.1.3. Characteristics and classification of plastic debris	10
2.1.3.1. Size classification of plastic debris	11
2.1.4. Plastic distribution, accumulation and enumeration	13
2.1.5. Degradation of plastic litter in marine environment	17
2.1.6. Impact on aquatic life	18
2.2. Persistent Organic Pollutants in Marine Plastic Debris	20
2.2.1. Persistent organic pollutants in marine environment	20
2.2.2. Passive samplers for persistent organic pollutants monitoring	20
2.2.3. Sorption and desorption of contaminants from plastics	22
2.2.4. Studies on persistent organic pollutants in beach plastics	23
2.2.5. Transfer of plastic-derived contaminants to organisms	26
2.3. Persistent Organic Pollutants in Sediment	27
2.3.1. Advances of analytical methods	29
2.3.2. Persistent organic pollutants in coastal sediment	31
2.3.3. Persistent organic pollutants in river sediment	33
2.3.4. Persistent organic pollutants in relation to sediment quality	33
2.3.5. Polycyclic aromatic hydrocarbons in sediment	36
2.3.6. Impact of persistent organic pollutants on organisms	39
2.3.7. Studies of persistent organic pollutants in India	41
3. MATERIALS AND METHODS	44
3.1 Study Area	44
3.1.1. Climate in Mumbai	45

3.2. Study Sites	45
3.2.1. Aksa beach	48
3.2.2. Versova beach	48
3.2.3. Juhu beach	48
3.2.4. Dadar beach	49
3.3. Sampling	49
3.3.1. Sampling for plastic litter in sediment	49
3.3.2. Plastic pellets collection for POP analysis	50
3.3.3. Sediment sampling for quality analysis	51
3.3.4. Sediment sampling for POP analysis	51
3.4. Sample Analysis	52
3.4.1. Sorting and quantification of plastic debris in sediment	52
3.4.2. Analysis of persistent organic pollutants in plastic pellets	52
3.4.2.1. Materials and reagents	52
3.4.2.2. Plastic pellets extraction in hexane	55
3.4.2.3. Fractionation of extract by silica gel column	57
3.4.2.4. Sample analysis by GC for PCB and pesticides	57
3.4.2.5. GC-MS analysis of PAHs	58
3.4.2.6. Identification and quantification of analytes in samples	58
3.4.2.7. Recovery estimation and method validation	61
3.4.3. Sediment quality analysis	62
3.4.3.1. pH	62
3.4.3.2. Electrical conductivity	62
3.4.3.3. Texture	62
3.4.3.4. Total organic carbon and total inorganic carbon	63
3.4.3.5. Available cations	63
3.4.4. Analysis of persistent organic pollutants in sediment	63
3.4.4.1. Soxhlet extraction	63
3.4.4.2. Cleanup and fractionation	64
3.4.4.3. Recovery estimation and method validation of POP analysis in sediment	64
3.5. Data Analysis	66

3.6. Statistical Analysis	67
4. RESULTS	68
4.1. Quantitative Assessment of Accumulation of Plastic Debris	68
4.1.1. Plastic litter enumeration in sediment	68
4.1.2. Size, colour and weight fractionation of plastic litter	73
4.1.2.1. Size fractionation of plastics	73
4.1.2.2. Correlation among abundances of size categories and total abundance	74
4.1.2.3. Colour fractionation of plastics	81
4.1.2.4. Weight fractionation of plastics	86
4.2. Persistent Organic Pollutants in Plastic Pellets	92
4.2.1. Polychlorinated biphenyls in plastic pellets	92
4.2.1.1. Spatial and temporal variation	93
4.2.1.2. Principal component analysis for PCBs	98
4.2.2. Organochlorine pesticides in plastic pellets	100
4.2.2.1. Spatial and temporal variation	101
4.2.2.2. Clustering of organochlorine pesticides	109
4.2.3. Polycyclic aromatic hydrocarbons in plastic pellets	109
4.2.3.1. Spatial and temporal variation	111
4.2.3.2. Polycyclic aromatic hydrocarbon distribution index	120
4.2.3.3. Clustering of polycyclic aromatic hydrocarbon compounds	122
4.3. Sediment Quality	122
4.4. Persistent Organic Pollutants in Sediment	126
4.4.1. Polychlorinated biphenyls in sediment	126
4.4.1.1. Spatial and temporal variation	127
4.4.1.2. Interaction of polychlorinated biphenyls and sediment properties	133
4.4.2. Organochlorine pesticides in sediment	136
4.4.2.1. Spatial and temporal variation	137
4.4.2.2. Interaction organochlorine pesticides and sediment properties	144
4.4.3. Polycyclic aromatic hydrocarbons in sediment	147
4.4.3.1. Spatial and temporal variation	148


4.4.3.2. Polycyclic aromatic hydrocarbon distribution index	157
4.4.3.3. Interaction of PAHs and sediment properties	158
4.5. Persistent Organic Pollutants in Plastic Pellets and Sediment	162
5. DISCUSSION	163
5.1. Contamination of Beach Sediment by Plastic Debris	163
5.1.1. Spatial and temporal variation	165
5.1.2. Size, color and weight fractionation	168
5.1.3. Mitigation of plastic pollution	173
5.2. Persistent Organic Pollutants in Plastic Pellets	174
5.2.1. Contamination of polychlorinated biphenyls in plastic pellets	177
5.2.2. Contamination of organochlorine pesticides in plastic pellets	183
5.2.3. Contamination of PAHs in plastic pellets	187
5.3. Persistent Organic Pollutants in Sediment	190
5.3.1. Contamination of polychlorinated biphenyls in sediment	191
5.3.2. Contamination of organochlorine pesticides in sediment	194
5.3.3. Contamination of polycyclic aromatic hydrocarbons in sediment	198
5.3.4. Environmental significance of POPs in sediment	202
5.4. Persistent Organic Pollutants in Plastic Pellets and Sediment	204
SUMMARY	207
REFERENCES	213

LIST OF TABLES		
Table No.	Title of the Table	Page No.
1	Field sampling programme from May 2011 to March 2012	50
2	Structure, chemical formula and molecular mass of 16 PAHs on the EPA list	54
3	IUPAC number and chemical names of the 7 PCBs	55
4	Chemical Structure and molecular mass of 16 OCPs	56
5	Retention times (min) of OCPs and PAHs standards	61
6	Summary of two-way ANOVA for total plastic debris (weight and number) among month, beach and their interactions	71
7	Summary of two-way ANOVA for size fractions of plastics among months, beaches and their interactions	75
8	Abundance of micro-plastics (mean±SE) by weight of particles (g m^{-2}) at beaches	77
9	Abundance of microplastics (mean±SE) by number of particles (items m^{-2}) at beaches	77
10	Abundance of macroplastics (mean±SE) by weight (g m^{-2}) at beaches	78
11	Pearson correlation matrix of weight and numbers of four size classes of plastics (n=72)	80
12	Summary of two-way ANOVA for distribution of colour fractions of plastics in sediment	82
13	Abundance of white plastics (mean±SE) by number (items m^{-2}) at beaches	83
14	Abundance of coloured plastics (mean±SE) by number (items m^{-2}) at beaches	84
15	Abundance of transparent plastics (mean±SE) by weight (g m^{-2}) at beaches	84

16	Summary of two-way ANOVA for distribution of weight fractions of plastics in beach sediment	88
17	Abundance of <0.05 g plastics (mean±SE) by weight (g m ⁻²) at beaches	89
18	Abundance of <0.05 g plastics (mean±SE) by number (items m ⁻²) at beaches	90
19	Abundance of 0.05-0.10 g plastics (mean±SE) by weight (g m ⁻²) at beaches	90
20	Mean and range of PCB compounds plastic pellets (n=72)	92
21	Summary of two-way ANOVA for distribution of ΣPCBs and PCB compounds in plastic pellets	95
22	Concentration of PCB compounds (mean±SE) at beaches during the study	97
23	Significant factor loadings (bold marked) for PCBs in plastic pellets	99
24	Mean±SE and range of 16 OCP compounds in plastic pellets (n=72)	101
25	Summary of two-way ANOVA for distribution of ΣOCPs, ΣHCHs and ΣDDTs in plastic pellets	102
26	Summary of two-way ANOVA for distribution of OCP compounds in plastic pellets	106
27	Concentration of OCP compounds (mean±SE) which varied significantly among beaches and/or months in plastic pellets	107
28	Mean±SE and range of PAH compounds, Σ16PAHs, ΣLPAHs and ΣHPAH, and potentially carcinogenic PAHs (ΣCPAHs) in plastic pellets	110
29	Summary of two-way ANOVA for distribution of ΣPAHs and PAH fractions in plastic pellets	112
30	Summary of two-way ANOVA for distribution of PAH compounds in plastic pellets	116

31	Bimonthly concentrations (ng g^{-1} ; mean \pm SE) of 2-3 ring compounds (mean \pm SE) in plastic pellets	117
32	Mean \pm SE and range of sediment quality parameters (n=72)	123
33	Summary of two-way ANOVA for distribution of pH, EC, TIC and TOC in sediment	123
34	Mean and range of seven PCB compounds in sediment (n=72)	126
35	Summary of two-way ANOVA for distribution of Σ PCBs and PCB compounds in sediment at $p=0.05$	129
36	Bimonthly and beach-wise concentrations (ng g^{-1} ; mean \pm SE) of PCB compounds in sediment	132
37	Pearson's multiple correlation matrix for 7 PCB compounds and some sediment quality parameters (n=72)	135
38	Mean \pm SE and range of 16 OCP compounds in beach sediment (n=72)	136
39	Summary of two-way ANOVA for distribution of Σ OCPs, Σ HCHs and Σ DDTs in sediment	139
40	Summary of two-way ANOVA for distribution of OCP compounds in sediment	142
41	Bimonthly concentrations (ng g^{-1} ; mean \pm SE) of OCP compounds in sediment	143
42	Pearson's multiple correlation matrix for Σ OCP, Σ HCH, Σ DDT, OCP compounds and some sediment quality parameters (n=72)	145
43	Mean \pm SE and range of 16PAH compounds, Σ PAHs, Σ LPAHs, Σ HPAH and potentially carcinogenic PAHs (Σ CPAHs) in sediment	148
44	Summary of two-way ANOVA for distribution of Σ PAHs and PAH fractions in sediment	150
45	Summary of two-way ANOVA for distribution of PAH compounds in sediment	153

46	Bimonthly and beach-wise mean concentration (ng g^{-1} ; mean \pm SE) of 2-3 ring PAH compounds in sediment	156
47	Pearson's multiple correlation matrix of mean Σ TPAHs, Σ LPAHs, Σ HPAHs, Σ CPAHs and sediment properties (n=72)	160
48	Pearson's multiple correlation matrix of mean PCB, OCP and PAH in plastic pellets and sediment	162
49	Abundances of microplastics in dry sediment in reported studies	171
50	Selected references reporting POP concentrations (ng g^{-1}) on plastic particles in marine environment	176
51	Selected references on concentrations of various POPs in sediment from different areas (ng g^{-1})	199
52	Comparison of OCPs and PCBs in the study area with ERL, ERM TEL and PEL values (ng g^{-1} dry wt.)	204



LIST OF FIGURES		
Figure No.	Title of the Figure	Page No.
1	The map showing the sampling sites in Mumbai coast	47
2	Plastic pellets sampled in Versova beach in July 2011	51
3	GC-ECD chromatograms of standard solutions (10 ng μl^{-1}): (a) PCBs, (b) OCPs	59
4	GC-MS chromatograms of standard solutions (10 ng μl^{-1}) of PAHs	60
5	Recovery (mean \pm SE) of PCBs in SRM-1944 (n=5)	65
6	Recovery (mean \pm SE) of OCPs in SRM-1944 (n=5)	65
7	Recovery of PAH compounds (mean \pm SE) in SRM-1944 (n=5)	66
8	Plastic litter abundance (mean \pm SE) in beach sediment of Aksa, Versova, Juhu and Dadar during the study period: (a) by weight, (b) by number of items	69
9	Microplastic items found on Juhu beach in May	70
10	Box-plots of plastic litter abundance in four beaches. The box represents 50% of the data in a series. The median is indicated by the horizontal line through the box. The ends of the box are the 25 th and 75 th percentiles. The error bars indicate the minimum and maximum observed values that are not statistically outlying. The extreme values are marked with an asterisk (*) in Juhu beach (n = 18 samples at each beach)	70
11	Abundance of plastics (mean \pm SE): (a) Bimonthly variation by weight and number; data points with different letters vary significantly among months in terms of number of plastics ($p < 0.05$). (b) Beach-wise variation by weight and number; bars with different letters in lowercase vary significantly among beaches for weight and uppercase for number of items ($p < 0.05$)	72

12	Abundance of micro-, meso-, macro- and mega-fractions (mean±SE) of plastics by weight and number: (a) Bimonthly variation (n=12), (b) Beach-wise variation (n=18)	76
13	Beach-wise variation of percentage of size fractions and overall mean (n=72) of plastic debris in sediment of Mumbai beaches by weight and number	78
14	Beach-wise variation of percentages of plastic pellets and overall mean (n=72) by number in microplastics (mean±SE)	79
15	Variation of colour fractions of plastics by weight and number: (a) Bimonthly variation (n=12), (b) Beach-wise variation (n=18)	81
16	Beach-wise percentages and overall mean (n=72) of colour fractions of plastic debris in beach sediment in terms of weight and number	85
17	Variation of weight fractions of plastics by weight and number: (a) Bimonthly variation (n=12), (b) Beach-wise variation (n=18)	87
18	Beach-wise percentages and overall mean (n=72) of weight fractions of plastic debris in terms of weight and number in beach sediment of Mumbai during the study period	91
19	Frequency of detection of seven PCB compounds in plastic pellets (n=72)	93
20	Bimonthly variation of Σ PCBs in Aksa, Versova, Juhu and Dadar beaches	94
21	Variation of Σ PCBs in plastic pellets: (a) Bimonthly variation (different letters within each month represent the significant difference at $p=0.05$), (b) Beach-wise variation	96
22	Beach-wise and overall mean of relative compositions of PCBs according to number of Cl atoms	98
23	Component plot in rotated space for PCBs in plastic pellets	99

24	Frequency of detection of 16 OCP compounds in plastic pellets (n=72)	100
25	Bimonthly variation of Σ OCPs in Aksa, Versova, Juhu and Dadar	102
26	Bimonthly variation of Σ HCHs and Σ DDTs in Aksa, Versova, Juhu and Dadar: (a) Σ HCHs, (b) Σ DDTs	103
27	Variation of Σ OCPs, Σ HCHs and Σ DDTs in plastic pellets: (a) Bimonthly variation (the values with different letters vary significantly at $p=0.05$ among months in terms of Σ DDTs), (b) Beach-wise variation (the values with different letters vary significantly at $p=0.05$ among beaches in terms of Σ DDTs)	104
28	Beach-wise variation and overall mean (n=72) of relative composition of HCHs, DDTs and other OCPs in plastic pellets	105
29	Dendrogram of 16 OCPs using complete lineage (Hierarchical cluster analysis)	109
30	Frequency of detection of 16 PAH compounds in plastic pellets (n=72)	111
31	Bimonthly variation of Σ PAHs in plastic pellets of Aksa, Versova, Juhu and Dadar	112
32	Variation of Σ PAHs, Σ LPAHs, Σ HPAHs, Σ CPAHs in plastic pellets: (a) Bimonthly variation (different letters in lowercase within each month represent the significant difference in terms Σ PAHs and different letters in uppercase within each month represent the significant difference in terms Σ LPAHs at $p=0.05$), (b) Beach-wise variation of (different letters within each site represent the significant difference in terms of Σ HPAHs at $p=0.05$)	113
33	Relative composition of LPAH and HPAH fractions in Σ PAHs in plastic pellets: (a) Bimonthly variation, (b) Beach-wise and overall means (n=72)	114
34	Beach-wise variation and overall means of carcinogenic PAHs percentage in plastic pellets	114

35	Beach-wise variation of 2-3 ring PAH compounds in plastic pellets (different letters in lowercase within each site represent the significant difference in terms of Ace and different letters in uppercase within each site represent the significant difference in terms of Phe at $p=0.05$)	115
36	Variation of 4 ring PAH compounds: (a) Bimonthly (different letters within each month represent the significant difference in terms of Pyr at $p=0.05$), (b) Beach-wise (different letters within each site represent the significant difference in terms of Flut at $p=0.05$)	118
37	Variation of 5-6 ring PAH compounds in plastic pellets: (a) Bimonthly (different letters within each month represent the significant difference in terms of B[a]P at $p=0.05$), (b) Beach-wise	119
38	Beach-wise variation and overall mean ($n=72$) of relative percentages of PAH compounds according to ring number	120
39	Variation of $\frac{\sum LPAHs}{\sum HPAHs}$ during the study: (a) Bimonthly (different letters within each month represent significant difference at $p=0.05$), (b) Beach-wise	121
40	Dendrogram based on complete linkage for PAH compounds in plastic pellets	122
41	Variation of pH and EC: (a) Bimonthly variation (different letters in lowercase within each site represent the significant difference in terms of pH and at $p=0.05$), (b) Beach-wise variation (different letters in uppercase within each site represent the significant difference in terms of EC at $p=0.05$)	124
42	Variation of inorganic and organic carbon: (a) Bimonthly variation (different letters in lowercase within each site represent the significant difference in terms of TIC at $p=0.05$), (b) Beach-wise variation (different letters in uppercase within each site represent the significant difference in terms of TOC at $p=0.05$)	125
43	Frequency of detection of 7 PCB compounds in sediment ($n=72$)	127

44	Bimonthly variation of Σ PCBs in sediment of Aksa, Versova, Juhu and Dadar	128
45	Variation of Σ PCBs from May 2011 to March 2012 in sediment: (a) Bimonthly (different letters within each month represent the significant difference at $p=0.05$), (b) Beach-wise (different letters within each month represent the significant difference at $p=0.05$)	130
46	Variation of Σ 3-5 Cl PCBs and Σ 6-7 Cl PCBs from May 2011 to March 2012 in sediment: (a) Bimonthly (different letters in lowercase within each month represent the significant difference for Σ 3-5 Cl PCBs and different letters in uppercase within each month represent the significant difference for Σ 6-7 Cl PCBs at $p=0.05$), (b) Beach-wise (different letters in lowercase within each beach represent the significant difference for Σ 3-5 Cl PCBs and different letters in uppercase within each beach represent the significant difference for Σ 6-7 Cl PCBs at $p=0.05$)	131
47	Beach-wise variation and overall mean ($n=72$) of relative compositions of PCB compounds in sediment	133
48	Dendrogram based on complete linkage for Σ PCBs, PCB compounds and sediment quality parameters	134
49	Frequency of detection of OCPs in sediment samples ($n=72$)	137
50	Bimonthly variation of Σ OCPs in sediment of Aksa, Versova, Juhu and Dadar	138
51	Bimonthly variation of Σ HCHs of sediment in Aksa, Versova, Juhu and Dadar	138
52	Bimonthly variation of Σ DDTs in sediment of Aksa, Versova, Juhu and Dadar	139
53	Variation of Σ OCPs, Σ HCHs and Σ DDTs in sediment: (a) Bimonthly, the values with different letters in lowercase vary significantly in terms of Σ OCPs, the values with different letters in lowercase (bold) vary significantly in terms of Σ HCHs, the values with different letters in uppercase vary significantly in terms of Σ DDTs at $p=0.05$; (b) Beach-wise	140

54	Beach-wise variation and overall mean (n=72) of relative composition of HCHs, DDTs and other OCPs in sediment	141
55	Dendrogram based on complete linkage for OCPs and sediment properties	144
56	Frequency of detection of 16 PAH compounds in sediment (n=72)	147
57	Bimonthly variation of Σ PAHs in sediment of Aksa, Versova, Juhu and Dadar beaches	149
58	Variation of Σ PAHs Σ LPAHs, Σ HPAHs, Σ CPAHs in sediment: (a) Bimonthly variation (different letters in lowercase, lowercase-bold and uppercase within each month represent the significant difference for Σ PAH, Σ LPAHs and Σ CPAHs, respectively, at $p=0.05$), (b) Beach-wise (different letters in lowercase within each site represent the significant difference in terms of Σ PAHs and different letters in uppercase within each site represent the significant difference in terms of Σ LPAHs at $p=0.05$)	151
59	Beach-wise variation and overall mean (n=72) of relative compositions of LPAHs and HPAHs in sediment	152
60	Variation of 4-ring PAH compounds in sediment: (a) Bimonthly (different letters in lowercase, uppercase and lowercase-bold within each month represent the significant difference for Flut, B[a]A and Chr, respectively, at $p=0.05$), (b) Beach-wise (different letters within each site represent the significant difference in terms of Chr at $p=0.05$)	154
61	Variation of 5-6 ring PAH compounds in sediment: (a) Bimonthly (different letters in lowercase and uppercase within each month represent the significant difference for I[cd]P and B[ghi]P respectively at $p=0.05$) (b) Beach-wise (different letters in lowercase and uppercase within each month represent the significant difference for I[cd]P and B[ghi]P respectively at $p=0.05$)	155

62	Beach-wise variation and overall mean (n=72) of relative composition of PAH compounds in sediment according to ring	157
63	Variation of $\frac{\sum LPAHs}{\sum HPAHs}$ in sediment during the study: (a) Bimonthly (different letters within each month represent the significant difference at $p=0.05$), (b) Beach-wise	158
64	Dendrogram based on complete linkage for ΣPAH , $\Sigma LPAH$, $\Sigma HPAH$ ΣCAH and PAH compounds, and some sediment quality parameters	159
65	Dendrogram based on complete linkage for POPs in plastic pellets and sediment	162